

Discovery of a New 89 Second X-ray Pulsar XTE J1906+09

D. Marsden, D. E. Gruber, W. A. Heindl, M. R. Pelling,
and R. E. Rothschild

Center for Astrophysics and Space Sciences, University of California at San Diego
La Jolla, CA 92093

Received _____; accepted _____

ABSTRACT

We report on the discovery of a new pulsating X-ray source during *Rossi X-ray Timing Explorer* observations of a low galactic latitude field centered at RA (J2000) = $19^{\text{hr}}05^{\text{m}}43^{\text{s}}$ and Dec (J2000) = $+08^{\circ}58'48''$. Significant pulsations were detected by both the PCA and HEXTE instruments aboard *RXTE* at a fundamental period of 89.17 ± 0.02 seconds, with higher harmonics also visible in the 2 – 10 keV power spectrum. The folded lightcurve from the source is multiply peaked at lower energies, and changes to single peaked morphology above ~ 20 keV. The phase averaged spectrum from the source is well fit by strongly absorbed power law or thermal bremsstrahlung spectral models of photon index 1.9 ± 0.1 or temperature 19.5 ± 4.6 keV, respectively. The mean neutral hydrogen column density is $N_H \approx 10^{23} \text{ cm}^{-2}$, suggesting a distance of > 10 kpc to the source and a minimum 2 – 10 keV X-ray luminosity of $2 \times 10^{35} \text{ ergs s}^{-1}$. By comparison with other pulsars with similar periods and luminosities, we suggest that XTE J1906+09 has a supergiant companion with an underfilled Roche lobe. We speculate further that one of the M stars in a peculiar M star binary system may be the companion.

Subject headings: pulsar: individual: XTE J1906+09 – stars: neutron – gamma-rays: bursts

1. Introduction

The discovery of a new member of a small class of astrophysical sources is important because each source has the potential to constrain theoretical models of the source physics. Neutron star binary systems are an example of a small class of X-ray sources, with numbers presently totaling total ~ 200 . Accretion-powered pulsars can be divided into two broad classes based on their X-ray luminosities and spectra (White, Swank, and Holt 1983). The X-ray spectra of the higher luminosity sources ($L_X \sim 10^{36} - 10^{37}$ ergs s $^{-1}$) are typically characterized by a hard power law of photon index $1 < \Gamma < 2$ out to an energy of 10 – 20 keV, above which the spectrum “breaks” to a much steeper index ($\Gamma > 3$). The lower luminosity accreting X-ray pulsars tend not to have this broken power law shape, and instead have a softer spectral shape ($\Gamma > 2$) over a broad range of X-ray energies. Furthermore, observations of low luminosity burst sources have revealed weak power law components ($\Gamma \sim 2$) when the inferred accretion accretion rate drops below a critical value (Barret & Vedrenne 1994). In this *Letter* we present the detection of a new source which appears to be a low luminosity accreting X-ray source.

2. Data

The 1° (FWHM) field of view centered at the point RA (J2000) = 286°.43 and DEC (J2000) = 8°.98 was observed by the High Energy X-ray Timing Experiment (HEXTE) and the Proportional Counter Array (PCA) instruments aboard the *Rossi X-ray Timing Explorer* (*RXTE*) satellite on a number of occasions between August 16-19, 1996, with a total livetime of ~ 25 ksec. The target of the observations was the Interplanetary Network (IPN) error box from the soft gamma-ray repeater SGR 1900+14 (Hurley et al. 1994), and the pointing was offset to avoid the bright confusing source 4U1907+097, which is $\sim 0^\circ.7$ away.

The HEXTE instrument consists of two clusters of collimated NaI/CsI phoswich detectors with a total net area of $\sim 1600 \text{ cm}^2$ and an effective energy range of $\sim 15 - 250 \text{ keV}$ (Rothschild et al. 1998). HEXTE background estimation utilizes near-real time observations of off-source viewing directions. The PCA instrument consists of 5 collimated Xenon proportional counter detectors with a total net area of 7000 cm^2 and an effective energy range of $2 - 60 \text{ keV}$ (Jahoda et al. 1996). For the PCA instrument, the instrumental background estimate is determined from modeling blank sky observations to estimate both the internal background of the detectors and the background due to cosmic X-ray flux.

3. Results

3.1. Timing Analysis

Photon event times from the PCA in the energy range $2 - 10 \text{ keV}$ were extracted and binned into lightcurves of 1 second binsize, after correcting the times to the solar system barycenter using the JPL DE200 ephemeris and *RXTE* pointing coordinates. The process was repeated for background data generated with the PCA background model, which was then subtracted from the on-source lightcurve. A Lomb-Scargle periodogram (Press & Rybicki 1989) was then formed from the data, resulting in the power spectrum shown in Figure 1 (top). Two harmonically spaced frequencies were detected at frequencies $f_2 \sim 0.0224 \text{ Hz}$ and $f_3 \sim 0.0336 \text{ Hz}$. The probability that each of these peaks is a random fluctuation is $\sim 10^{-21}$, given the number of frequencies examined and the exponential probability distribution of the Lomb-Scargle periodogram. The presence of peaks corresponding to f_5 and f_6 is also suggested by the periodogram, but at a marginal significance. Periodogram and FFT analysis using different time binsizes, no background subtraction, individual PCA detector units, and time interval subsets all show evidence for of one or two harmonic peaks in the power spectra.

The harmonically-spaced frequencies f_2 and f_3 imply a fundamental frequency of $f_1 \sim 0.0112$, corresponding to a period of ~ 89 seconds. To search for the fundamental period we performed a chi-squared folding analysis on 128 frequencies corresponding to periods about 89 seconds, with a frequency resolution equal to the independent Fourier spacing implied by the total duration of the data. The result, shown in Figure 1 (bottom), indicates a fundamental period of 89.17 ± 0.015 seconds. We then searched the HEXTE event data for this periodicity, using the Z_n^2 test (Buccheri et al. 1983). HEXTE on-source event times in the energy range 15 – 50 keV were accumulated and corrected to the solar system barycenter using the the JPL DE200 ephemeris and *RXTE* pointing coordinates (as before). The 4×10^5 event times yielded a Z_1^2 value of 75.3, using the single frequency corresponding to the period 89.17 seconds. This corresponds to a random probability of 5×10^{-17} , given the probability distribution of Z_n^2 . As a check, the 15 – 50 keV HEXTE *off-source* events were analyzed in an identical manner and a value of $Z_1^2 = 0.3$ was obtained — consistent with no periodicity.

The barycenter-corrected PCA and HEXTE event times were extracted for various energy ranges and folded on the fundamental period $P = 89.17$ seconds (Figure 2). The corrected event times were normalized to MJD 50342.0 (TDB) before folding. The folded lightcurve (FLC) appears to change at energies between 20 – 30 keV, with the pulse profile becoming more singly-peaked and possibly shifting in phase. In addition, the pulsed emission for energies above ~ 20 keV peaks at a different phase than the emission at lower energies. Given the contributions from instrumental, cosmic X-ray, and Galactic ridge fluxes implied by the spectral fits (discussed below), we derive pulsed fractions of 16%, 6%, and $\sim 4\%$ in the energy ranges 2 – 10, 10 – 20 and 20 – 200 keV, respectively.

We have searched for evidence of Doppler shifts associated with possible binary motion in the observed periodicities from XTE J1906+09. Barycenter corrected 2 – 60 keV PCA

event times were collected for 7 observation intervals throughout the observing period. The frequencies f_2 and f_3 were determined for each time interval via fitting of the peak in the Z_1^2 distribution, taken from small frequency ranges about $f_2 = 0.0224$ and $f_3 = 0.0336$, with a Gaussian. The fundamental frequency f_1 (corresponding to the 89.17 s period) was then obtained for each data set by the relation $f_1 = \frac{1}{5}(f_2 + f_3)$. We find that the fundamental frequency obtained in this manner is consistent with a constant frequency over the 3 day observation period. Because of the limited (7) number of observations and limited temporal coverage of the observations, however, the non-detection of significant Doppler shifts poses no serious constraint on the possible binary orbital parameters of the system. For secular period changes, we fit a model to the data in which the period changes linearly with time, and obtained a 2σ upper limit of $|\dot{P}| < 2 \times 10^{-6}$.

3.2. Spectral Analysis

Spectral analysis of relatively faint X-ray sources at low Galactic latitudes is complicated by the presence of diffuse emission from the Galactic ridge. Recent work by Valinia & Marshall (1998) indicates that the pointing direction of XTE J1906+09 is outside the bright, thin component of the ridge emission but inside the broader (FWHM $\sim 4^\circ$) region. Only one of the HEXTE background positions is centered on a region of significantly higher Galactic X-ray flux $(l, b) = (39.6^\circ, 0.3^\circ)$, and this background is excluded from the analysis. Following Valinia & Marshall (1998), we model the Galactic ridge emission in the PCA with thermal and nonthermal components consisting of Raymond-Smith plasma (with solar elemental abundances and zero redshift) and power law spectral shapes, respectively, multiplied by a neutral hydrogen absorption factor.

The phase-averaged PCA and HEXTE data were fit simultaneously to various continuum spectral models using XSPEC 10.0 and assuming that XTE J1906+09 was at

the center of the field of view. For the spectral fits, PCA and HEXTE data in the energy ranges $2.5 - 30$ and $15 - 250$ keV, respectively, were used. To account for uncertainties in the effective open area of the two instruments, the relative normalization of HEXTE was treated as a free parameter in the spectral fits. The plasma temperature, spectral index, and N_H values of the Galactic ridge spectrum (PCA only) were held fixed at values appropriate for the “Central Ridge” (see Table 3 from Valinia & Marshall 1998), but the normalizations were allowed to vary, to account for the fact that the *RXTE* pointing position is displaced from the Galactic plane by 1° . The results for a power law model for the XTE J1906+09 emission is given in Table 1. A thermal bremsstrahlung spectral shape of temperature 19.5 ± 4.6 keV and $N_H = (7.7 \pm 5.2) \times 10^{22} \text{ cm}^{-2}$ fits the data equally well. More complicated spectral models (such as broken power laws) were not fit to the data, due to insufficient counts above ~ 20 keV. The reduced chi-squared values for both the power law and thermal bremsstrahlung spectral fits were 1.0 for 523 degrees of freedom. As expected, for both fits the Galactic ridge normalization values were lower than the values given in Valinia & Marshall (1998) — consistent with the 1° offset from the Galactic plane — and the relative normalization factor between the PCA and HEXTE was consistent with previous fits to high Galactic latitude sources (Marsden et al. 1997). In the spectral fitting, we have assumed that all the observed iron line emission is from the Galactic ridge (Raymond-Smith plasma component), and therefore we do not attempt to constrain the iron line emission from XTE J1906+09.

4. Discussion

Because of the low galactic latitude of the observed field, there is no shortage of possible XTE J1906+09 counterparts. The *ROSAT* all-sky survey (Voges et al. 1996) sources RX J190724+0843.4 and RX J190717+0919.3 (Vasisht, Kulkarni, Frail, & Greiner 1994) are

unlikely counterparts, because their count rates are significantly higher (by at least an order of magnitude) than the predicted *ROSAT* PSPC count rate of XTE J1906+09, given the spectral parameters listed in Table 1. The predicted *ROSAT* HRI count rates are 7×10^{-5} and 2.8×10^{-4} counts s^{-1} for the power law and thermal bremsstrahlung spectral forms, respectively. Supernova remnants such as G42.8+0.6 and other supernova remnants in the field of view (Vasisht, Kulkarni, Frail, & Greiner 1994) are usually not associated with slow (period > 10 s) pulsars because of age constraints.

The pulsation period and inferred luminosity of XTE J1906+09 indicate that the X-ray emission is powered by accretion onto a neutron star. The high absorption column density implied by the spectral fit suggests that the source is located at a distance of at least ~ 10 kpc, and this yields a lower limit to the X-ray luminosity ($2 - 10$ keV) of 2×10^{35} ergs s^{-1} . A variable pulse shape with energy (Fig. 2) is characteristic of accretion-powered neutron star systems, in which the emergent X-ray emission pattern is determined by the geometry and the distribution of matter in the vicinity. The scattering cross section is also energy dependent, and hence the pulse profile is expected to vary with energy. In addition, at lower energies more complicated pulsar lightcurves are expected due to photoelectric absorption by material around the neutron star.

Assuming that XTE J1906+09 is an accretion-powered binary system, we can speculate on the nature of the system. The low luminosity and long spin period of XTE J1906+09 are atypical of low mass X-ray binary systems, but are consistent with a high mass system in which the accretion takes place via mass ejection from the companion star. Such high mass X-ray binaries can be divided into two classes: underfilled Roche lobe systems and Be-binary systems. Be-binary systems are transient systems in which the outflow from the Be star companion takes the form of either a stellar wind or an equatorial mass ejection (Rappaport, S. & Van den Heuvel 1982). These sources are transient or highly variable,

with X-ray luminosities ranging over ~ 4 orders of magnitude ($L_X \sim 10^{34} - 10^{38}$). If XTE J1906+09 is such a system, it probably has a long orbit ($P_{orb} > 50$ days), due to the observed correlation between orbit and pulsar periods in Be-binaries (Bildsten et al. 1997). One problem with this identification is that the X-ray spectra of these systems are characterized by power laws with photon indices > 3 ($\sim 20 - 100$ keV, Bildsten et al. 1997). The $\sim 20 - 250$ keV spectrum of XTE J1906+09 (using the HEXTE data only) is fit by a power law of photon index 1.45 ± 0.45 , which is significantly harder than this. In addition, there are no known Be stars in the 1° XTE J1906+09 error box (Jaschek & Egret 1982).

XTE J1906+09 could also be an underfilled Roche lobe supergiant system. These systems typically have long orbital periods ($P_{orb} > 2$ days), long spin periods ($P_{spin} > 100$ s), and relatively low X-ray luminosities of $10^{35} - 10^{37}$ ergs s $^{-1}$ (Bildsten et al. 1997). The inferred parameters of XTE J1906+09 are consistent with being a member of this class, but the X-ray spectral index of XTE J1906+09 is flatter than that seen from Roche lobe supergiant systems (Bildsten et al. 1997).

The peculiar double M star infrared system discovered by Hartmann et al. (1995) is also in the *RXTE* field of view. The stars in this system are heavily reddened ($A_V \approx 19.2$), implying a distance of $12 - 15$ kpc, and are thought to be gravitationally bound with an orbit of 2×10^6 years (Vrba et al. 1996). Using the average extinction curve given in Savage & Mathis (1979), and the relation between N_H and $E(B - V)$ given in Korneef (1982), we derive the relation $A_V \approx 1.6N_{H22}$ mag, where N_{H22} is N_H in units of 10^{22} cm $^{-2}$. From the mean XTE J1906+09 N_H value obtained from the power law and bremsstrahlung fits, we obtain $A_V = 15 \pm 5$ mag, which is consistent with the value seen from these stars.

Given the similarities between the observed properties of XTE J1906+09 and high mass wind-fed X-ray binary systems, and the similar inferred distances of XTE J1906+09 and the M star supergiant system, we regard the latter as a promising counterpart candidate to

XTE J1906+09. The large separation of the M stars would allow one to have a neutron star companion without significant effect on the M star system. Then the M star - neutron star system could be the source of the X-ray flux seen by *RXTE*. Vrba et al. (1996) have suggested that such a combination might be the counterpart of the soft gamma-ray repeater SGR 1900+14. The detection of 89 second pulsations from these IR sources would verify the association of XTE J1906+09 with this system, and XTE J1906+09 would become the first known X-ray triple system.

5. Conclusions

We have presented the detection of a new 89 second pulsating X-ray source with *RXTE*. The source was detected during observations of a region of Galactic plane containing the SGR 1900+14 error box. Characteristics of the folded lightcurves and spectrum suggest that the source, XTE J1906+09, is a low luminosity X-ray binary located beyond the Galactic center. The low luminosity and long pulse period of XTE J1906+09 indicate that this source is probably a high mass X-ray binary accreting via a stellar wind. We raise the possibility that the new source is associated with the highly absorbed double M supergiant binary system on the edge of the SGR 1900+14 error box. If this identification is correct, XTE J1906+09 is the first known X-ray triple system, and a candidate counterpart for SGR 1900+14.

We would like to thank the duty scientists at the online *RXTE* help desk for assistance in analyzing the PCA data. This work was funded by NASA grant NAS5-30720.

REFERENCES

- Barret, D. & Vedrenne, G. 1994, ApJS, 92, 505
- Bildsten, L. et al. 1997, ApJS, 113, 367
- Buccheri, R. et al. 1983, A&A, 128, 245
- Córdova, F. A. 1995, in *X-ray Binaries*, ed. W. H. G. Lewin, J. van Paradijs, & E. P. J. van den Heuvel (Cambridge: Cambridge University Press), 352
- Hartmann, D. H. et al. 1995, in AIP conf. Proc. 366, Workshop on High Velocity Neutron Stars, ed. R. E. Rothschild & R. E. Lingenfelter (New York: AIP), 84
- Hurley, K. et al. 1994, ApJ, 431, L31
- Jahoda, K. et al. 1996, EUV, X-ray, and Gamma-Ray Instrumentation for Astronomy VII, SPIE Proceedings, eds: O. H. V. Sigmund and M. Gumm, 2808, 59
- Jaschek, M. and Egret, D. 1982, IAU Symposium No. 98, “Be Stars”, eds. Jaschek, M. & Groth, H.G., 261
- Korneef, J. 1982, A&A, 107, 247
- Marsden, D. et al. 1997, ApJ, 491, L39
- Press, W. H. & Rybicki, G. B. 1989, ApJ, 338, 277
- Rappaport, S. & Van den Heuvel, E.P.J. 1982, IAU Symposium No. 98, “Be Stars”, eds. Jaschek, M. & Groth, H.G., 327
- Rothschild, R. E. et al. 1998, ApJ, 496, 538
- Savage, B. D. & Mathis, J. S. 1979, ARA&A, 17, 73

Valinia, A. & Marshall, F. E. 1998, ApJ, in press

Vasisht, G., Kulkarni, S. K., Frail, D. A. & Greiner, J. 1994, ApJ, 431, L35

Voges, W. et al. 1996, IAU Circular, 6420

Vrba, R. J. et al. 1996, ApJ, 468, 225

White, N. E., Swank, J. H., and Holt, S. S. 1983, ApJ, 270, 711

Fig. 1.— The detection of periodicity. The top panel shows the Lomb periodogram of 2–10 keV binned time series data from the PCA, with the harmonics labeled. The bottom panel shows the results of a chi-squared folding search for the fundamental period (89 seconds) of the pulsations, using the same data.

Fig. 2.— The folded lightcurve of the 89 second pulsations over a range of energies. The two middle panels correspond to the same energy range (see text). The background has not been subtracted.

Table 1. Phase-Averaged Spectral Fit Results

Model Parameter	XTE J1906+09	Galactic Ridge
N_H^a	10.8 ± 4.0	1.8 (fixed)
Photon Index	1.9 ± 0.1	1.8 (fixed)
Power Law Flux ^b	16.0 ± 2.5	9.0 ± 0.4
Plasma Temperature ^c	–	2.9 (fixed)
Plasma Flux ^b	–	7.7 ± 0.7

^aNeutral hydrogen absorption (10^{22} H atoms cm^{-2})

^bUnabsorbed 2 – 10 keV flux (10^{-12} ergs cm^{-2} s^{-1})

^cRaymond-Smith plasma temperature (keV)



